

The Nordic Liver Transplant Registry (NLTR)

Annual report 2015

Report prepared by Espen Melum June 2016

Responsible contact persons:

Scandiatransplant Denmark - Århus; Ilse Duus Weinreich

Denmark - Copenhagen; Allan Rasmussen

Sweden - Gothenburg; William Bennet

Sweden - Stockholm; Bo-Göran Ericzon

Finland - Helsinki; Helena Isoniemi

Norway - Oslo; Bjarte Fosby

NLTR; Espen Melum espen.melum@medisin.uio.no

1. Source of data

The numbers and graphs included in the present report are based on data extracted from the Nordic Liver Transplant Registry (NLTR) in April 2016. Prior to this export, data were subjected to extensive integrity and quality control. Entry of missing data and correction of errors were performed meticulously by transplant coordinators at all centers prior to the final data extraction.

2. Data content NLTR 2015

The registry comprises complete data from the liver transplantation activity at all transplantation centres in Denmark, Sweden, Norway and Finland since 1982. Before 1990, only patients that were transplanted were registered. After 1990, the registry covers all patients entered to the liver transplantation waiting list, regardless of transplantation status. From September 1994, complete waiting list data are available from all patients in addition to the transplantation details. Data are stored securely at Scandiatransplant in Århus (www.scandiatransplant.org).

Up to December 31st 2015, data from a total of 6820 patients had been entered to the NLTR. Of these, 5913 patients had received a first liver graft. Of these, 589 (10.0%) had been transplanted more than once, and 87 (1.5%) had been transplanted more than twice. A total of 166 living donor transplantations had been performed. Children below 16 years constituted 610 (10.3%) of the transplanted patients in the registry.

3. Transplantation activity 2015

The total number of patients who received a first liver graft in 2015 was 362 (Figure 1). Of these, 4 were combined liver-kidney transplantations. Of the first liver transplantations in 2015 5 were

living donor transplantations and 2 domino transplantations. Three of the living donor transplantations were performed in Oslo. In addition, 39 re-transplantations were performed (Table 2). The total number of 401 liver transplantations represents an increase from the 388 liver transplantations performed in 2014 (Figure 1) and the number of liver transplantations in the Nordic countries is for the first time over 400. The number of re-transplantations remains steady (Figure 1).




Figure 1. Number of patients receiving a liver allograft 1982-2015. The blue line represents the number of patients receiving a first liver graft while the red line represents the total number of re-transplantations.

	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Copenhagen	32	37	43	37	43	42	44	39	41	55
Gothenburg	52	64	66	78	61	67	75	72	89	86
Helsinki	49	50	42	42	47	52	48	44	56	70
Oslo	52	64	69	69	77	81	89	96	89	72
Stockholm	56	50	52	43	53	65	69	73	77	79
Uppsala	8	8	11	10	3	0	0	0	0	0

Table 1. Number of first liver transplantations performed at the individual centers during the last 10 years.

	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Copenhagen	4	5	1	3	4	9	4	3	6	3
Gothenburg	8	11	10	11	19	16	4	9	8	7
Helsinki	4	3	5	6	3	4	4	5	3	7
Oslo	10	8	10	13	12	8	11	14	11	12
Stockholm	3	2	6	3	1	8	5	7	8	6
Uppsala	0	1	2	1	0	0	0	0	0	0

Table 2. Total number of re-transplantations performed at the individual centers during the last 10 years.




Figure 2. Number of first liver transplantations performed at Nordic centers that are currently performing liver transplantations.




Figure 3. Number of first liver transplantations performed in the Nordic countries according to the country's population. PMP, per million population.

4. The waiting list 2015

In 2015, a total of 381 patients were entered into the waiting list for a first liver transplant (Table 3) which is a decline from the 417 entered in 2014 (Figure 4). Twenty-four of the patients entered in 2015 were listed as highly urgent.

Active on waiting list	Deceased donor	Living donor	Dead	Permanent withdrawal
83	273	7	10	6

Table 3. Patients entering the waiting list in 2015 classified by outcome as of December 31st 2015.




Figure 4. Number of patients entering the waiting list from 2000-2015.

The number of deaths among patients waiting for a first liver transplant in 2015 was 10 (Denmark 2, Sweden 5, Finland 1, Norway 2). The absolute number of deaths registered on the waiting list has remained stable since 1990 (Figure 5). When the deaths on the waiting list are evaluated in relation to the total liver transplantation activity there has been a decrease in the number of deaths on the waiting list (Figure 6).




Figure 5. Number of patients registered as dead on the waiting list in the period 1990-2015.




Figure 6. Number of patients registered as dead on the waiting list relative to the total transplantation activity in the period 1990-2015.

The median waiting time in 2015 was 39 days when excluding patients listed for a highly urgent liver transplantation. The differences according to different ABO blood types were as expected (Table 4) with largely similar numbers since 2000 (Figure 7).

0	A	AB	B
66 (715)	32 (661)	26 (254)	39.5 (318)

Table 4. Median time on waiting list (days) for patients receiving a first liver allograft in 2015 according to ABO blood type. The number in parenthesis represents the maximum waiting time for the indicated blood type in 2015. (Patients listed as highly urgent are excluded from the calculations.)




Figure 7. Median waiting time for first liver transplantation according to ABO blood type for 2000-2015. (Patients listed as highly urgent are excluded from the calculations.)

There were slight differences in the median waiting time between the centers in 2015 (Table 5). There has been a steep decrease in the median waiting time for Danish patients over the last years and now it

seems to have stabilized on the same level as the other centers. A slight decrease in the median waiting time is seen in Norway and Finland (Figure 8).

Copenhagen	Gothenburg	Helsinki	Oslo	Stockholm
56 (343)	47.5 (661)	28.5 (333)	26 (715)	53.5 (420)

Table 5. Median time on waiting list (days) for patients receiving a first liver allograft in 2015 according to transplantation center. The number in parenthesis represents the maximum waiting time for the indicated center in 2015. (Patients listed as highly urgent are excluded from the calculations.)




Figure 8. Median waiting time for first liver transplantation according to country for 2000-2015. (Patients listed as highly urgent are excluded from the calculations.)

5. Age of recipients and donors

The mean age of adult liver recipients (≥ 16 years, first liver transplantation) in 2015 was 53.4 years. Mean age of children (< 16 years, first liver transplantation) in 2015 was 6.6 years. Since 1990 the proportion of recipients > 60 years of age at the first transplantation has gradually increased (Figure 9). The mean age of the donors has increased since 1990 with a similar trend in all the Nordic countries (Figure 10).




Figure 9. Proportion of liver transplants in the indicated age groups.




Figure 10. Mean age of donors utilized in the indicated years stratified for the different Nordic countries.

6. Diagnoses

In 2015, primary sclerosing cholangitis and hepatocellular carcinoma were the leading indications for liver transplantation in the Nordic countries (Table 6). Of the patients listed for transplantation with a primary diagnosis of hepatocellular carcinoma (HCC) in 2015 36% were also anti-HCV positive.

	1982-90	1991-95	1996-00	2001-05	2006-10	2011-15	2015
Primary sclerosing cholangitis	10.9%	12.2%	15.6%	15.6%	15.7%	16.6%	19.7%
Hepatocellular carcinoma	10.9%	4.3%	6.0%	6.6%	10.4%	17.1%	17.3%
Alcoholic cirrhosis	1.9%	8.3%	11.6%	12.8%	10.9%	12.1%	13.6%
Metabolic disease	10.0%	8.1%	6.0%	4.8%	7.0%	7.0%	8.1%
Cirrhosis - unknown	0.6%	2.6%	3.5%	2.4%	5.0%	6.7%	7.1%
Acute liver failure - other	8.4%	10.7%	7.4%	6.9%	5.9%	5.2%	6.0%
Primary biliary cirrhosis	22.2%	14.5%	8.9%	6.7%	6.7%	5.2%	4.7%
Autoimmune cirrhosis	2.8%	3.5%	3.5%	4.3%	4.4%	4.6%	4.5%
Post hepatitis C cirrhosis	0.0%	3.4%	8.1%	10.1%	10.5%	7.1%	3.4%
Polycystic disease	0.3%	1.0%	1.2%	1.4%	1.6%	1.8%	2.6%
Biliary atresia	6.9%	5.6%	4.4%	4.8%	2.5%	2.2%	1.8%
Acute liver failure - toxic	0.6%	2.7%	3.9%	4.3%	3.8%	2.6%	1.6%
Post hepatitis B cirrhosis	0.9%	2.7%	3.1%	3.3%	1.6%	1.2%	1.3%
Other liver malignancies	2.5%	2.3%	2.1%	1.5%	0.9%	1.5%	1.0%
Secondary liver tumors	0.9%	0.6%	0.6%	0.5%	2.2%	1.9%	1.0%
Biliary tract carcinoma	0.0%	0.0%	0.0%	0.1%	0.7%	0.5%	0.8%
Post hepatitis D cirrhosis	0.0%	0.0%	0.0%	0.0%	0.2%	0.6%	0.8%
Secondary biliary cirrhosis	0.9%	0.5%	0.2%	0.2%	0.3%	0.4%	0.8%
Acute liver failure - viral	1.3%	2.7%	1.6%	1.0%	1.7%	1.3%	0.5%
Cholangiocarcinoma	1.6%	0.9%	0.8%	0.7%	0.9%	0.1%	0.5%
Others	16.5%	13.4%	11.3%	12.0%	7.4%	4.2%	2.7%

Table 6. Diagnoses of patients listed for a first liver transplantation in 2015 compared with previous years. In 2015 36% of HCC patients listed for a first liver transplantation were anti-HCV positive.

7. Patient and liver graft survival

When looking at 5-years intervals, patient survival (defined as time from the first liver transplantation until death) and graft survival (defined as time from the first liver transplantation until death or re-transplantation) were dramatically improving over the first years of the Nordic liver transplantation programs (Figures 10 and 11). It is now evident that there is a further increase in the observed survival also in the most recent 5-year period. There are notable differences in the long-term patient and graft survival for different indications for transplantation (Figures 12 and 13).




Figure 10. Kaplan-Meier patient survival curve for patients receiving a first liver allograft in the indicated time periods.




Figure 11. Kaplan-Meier graft survival curve for patients receiving a first liver allograft in the indicated time periods.




Figure 12. Kaplan-Meier patient survival curve for patients receiving a first liver allograft stratified for the three most common primary diagnoses.




Figure 13. Kaplan-Meier graft survival curve for patients receiving a first liver allograft stratified for the three most common primary diagnoses.

8. Maintenance of the registry

There are notable differences between each center in terms of how extensively data are entered into the NLTR. Diagnosis information, waiting list/transplantation status and survival data for all patients are now complete for 2015. I am extremely grateful for the meticulous follow-up provided by the transplant coordinators upon my requests during quality control. In Oslo, I particularly want to thank Stein Foss, in Gothenburg Christina Wibeck and Ulla Nyström, in Stockholm Marie E. Larsson, in Copenhagen Mette Gottlieb and in Helsinki, Helena Isoniemi is overseeing the maintenance of the registry. Quality control of the content of NLTR is a continuous priority, and a particular emphasis is put into ensuring integrity of the survival data,

including cause of death. The remainder of the registry must be maintained at a level set at the discretion of each individual center and contact person. Data transfer between NLTR and the ELTR are in the process of being established. In 2015 the interface for NLTR was transferred to the new web-based system YASWA together with the rest of Scandiatransplant. This process went extremely smooth thanks to good planning from Scandiatransplant.

9. Acknowledgements - financial support

The NLTR received no financial support in 2015. The maintenance of the Oracle system has been performed by Scandiatransplant. We are extremely grateful for the help and support from Frank Pedersen and Ilse Duus Weinreich and the rest of the Scandiatransplant team in Aarhus. Without their assistance it would very simply not have been possible to maintain the registry and I sincerely hope their efforts are recognized by the NLTG and Scandiatransplant.

10. Organization and data ownership

The registry (software) is the property of Scandiatransplant. The data in the registry are the property of the hospitals represented in the Nordic Liver Transplantation Group. Utilization of data in research projects should be censored by the latter and need to comply with national guidelines for research ethics and data handling. Co-authorships for publications from research projects should be allocated according to the Vancouver guidelines, this includes presentations of data at conferences. The quality statistics of the transplantation activity presented in this report must not be used in other contexts without permission from the Nordic Liver Transplantation Group.

11. Publications based on the NLTR

Full length articles 1990-2015:

1. Keiding S, Ericzon BG, Eriksson S, Flatmark A, Hockerstedt K, Isoniemi H, Karlberg I, Keiding N, Olsson R, Samela K, Schrumpf E. Survival after liver transplantation of patients with primary biliary cirrhosis in the Nordic countries. Comparison with expected survival in another series of transplantations and in an international trial of medical treatment. *Scand J Gastroenterol* 1990; 25:11-8
2. Hockerstedt K, Ericzon BG, Eriksson LS, Flatmark A, Isoniemi H, Karlberg I, Keiding N, Keiding S, Olsson R, Samela K. Survival after liver transplantation for primary biliary cirrhosis: use of prognostic indices for comparison with medical treatment. *Transpl Proc* 1990; 22:1499-500
3. Hockerstedt K, Isoniemi H, Ericzon BG, Broome U, Friman S, Persson H, Bergan A, Schrumpf E, Kirkegaard P, Hjortrup A. Is a 3-day waiting list appropriate for patients with acute liver failure? *Transpl Proc* 1994;26:1786-7
4. Bjøro K, Friman S, Höckerstedt K, Kirkegaard P, Keiding S, Schrumpf E, Olausson M, Oksanen A, Isoniemi H, Hjortrup A, Bergan A, Ericzon BG. Liver transplantation in the Nordic countries, 1982-1998: Changes of indications and improving results. *Scand J Gastroenterol* 1999;34:714-722
5. Bjøro K, Höckerstedt K, Ericzon BG, Friman S, Hjortrup A, Keiding S, Schrumpf E, Duraj F, Olausson M, Mäkisalo H, Bergan A, Kirkegaard P. Liver transplantation in patients over 60 years of age. *Transpl Int* 2000; 13, 165-170
6. Bjøro K, Kirkegaard P, Ericzon BG, Friman S, Schrumpf E, Isoniemi H, Herlenius G, Olausson M, Rasmussen A, Foss A, Höckerstedt K. Is a 3-day limit for highly urgent liver transplantation for fulminant hepatic failure appropriate – or is the diagnosis in some cases incorrect? *Transpl Proceed* 2001;33:2511-3

7. Ericzon BG, Bjøro K, Höckerstedt K, Hansen B, Olausson M, Isoniemi H, Kirkegaard P, Broome U, Foss A, Friman S. Time to request ABO-identity when transplanting for fulminant hepatic failure? *Transpl Proc* 2001;33:3466-7
8. Leidenius M, Broome U, Ericzon B-E, Friman S, Olausson M, Schrumpf E, Höckerstedt K. Hepatobiliary carcinoma in primary sclerosing cholangitis: a case control study. *J Hepatol* 2001; 34: 792-8.
9. Olausson M, Mjornstedt L, Backman L, Lindner P, Olsson R, Krantz M, Karlens KL, Stenqvist O, Henriksson BA, Friman S. Liver transplantation--from experiment to routine care. Experiences from the first 500 liver transplantations in Gothenburg. *Lakartidningen* 2001;98:4556-62
10. Brandsæter B, Höckerstedt K, Ericzon BG, Friman S, Kirkegaard H, Isoniemi H, Foss A, Olausson M, Hansen B, Bjøro K: Outcome following listing for liver transplantation due to fulminant hepatic failure in the Nordic countries. *Liver Transplantation* 2002;8:1055-62
11. Bjøro K, Ericzon BG, Kirkegaard P, Höckerstedt K, Söderdahl G, Olausson M, Foss A, Schmidt LE, Brandsæter B, Friman S. Liver transplantation for fulminant hepatic failure: impact of donor-recipient ABO-matching on the outcome. *Transplantation* 2003; 75:347-53
12. Brandsæter Bjørn, Broomé Ulrika, Isoniemi Helena, Friman Styrbjörn, Hansen Bent, Schrumpf Erik, Oksanen Antti, Ericzon Bo-Göran, Höckerstedt Krister, Mäkisalo Heikki, Olsson Rolf, Olausson Michael, Kirkegaard Preben, Bjøro Kristian. Liver transplantation for primary sclerosing cholangitis in the Nordic countries: outcome after acceptance to the waiting list. *Liver Transpl.* 2003;9:961-9.
13. Brandsæter B, Friman S, Broome U, Isoniemi H, Olausson M, Backman L, Hansen B, Schrumpf E, Oksanen A, Ericzon BG, Höckerstedt K, Mäkisalo H, Kirkegaard P, Bjøro K. Outcome

following liver transplantation for primary sclerosing cholangitis in the Nordic countries. *Scand J Gastroenterol.* 2003;38:1176-83.

14. Brandsaeter B, Isoniemi H, Broome U, Olausson M, Backman L, Hansen B, Schrumpf E, Oksanen A, Ericzon BG, Hockerstedt K, Makisalo H, Kirkegaard P, Friman S, Bjoro K. Liver transplantation for primary sclerosing cholangitis; predictors and consequences of hepatobiliary malignancy. *J Hepatol.* 2004;40:815-822.

15. Bjoro K, Schrumpf E. Liver transplantation for primary sclerosing cholangitis. *J Hepatol.* 2004;40:570-7.

16. Brandsaeter B, Isoniemi H, Broomé U, Olausson M, Bäckmann L, Hansen B, Oksanen A, Ericzon BG, Höckerstedt K, Mäkisalo H, Kirkegaard P, Friman S, Bjoro K, Schrumpf E (Nordic Liver Transplantation Group). Chemopreventive effect of ursodeoxycholic acid in primary sclerosing cholangitis? Falk Symposium 141. *Bile Acid Biology and its Therapeutic Implications. XVIII International Bile Acid Meeting (2005; page 242-249).*

17. Melum E, Schrumpf E, Bjoro K. Liver TX for hepatitis C cirrhosis in a low prevalence population: risk factors and status at evaluation. *Scand J Gastroenterol.* 2006;41:592-6.

18. Bjoro K, Brandsaeter B, Foss A, Schrumpf E. Liver transplantation in primary sclerosing cholangitis. *Semin Liver Dis.* 2006;26:69-79.

19. Melum E, Friman S, Bjoro K, Rasmussen A, Isoniemi H, Gjertsen H, Bäckman L, Oksanen A, Olausson M, Duraj FF, Ericzon BG. Hepatitis C impairs survival following liver transplantation irrespective of concomitant hepatocellular carcinoma. *J Hepatol.* 2007 Dec;47(6):777-83.

20. Friman S, Foss A, Isoniemi H, Olausson M, Höckerstedt K, Yamamoto S, Karlsen TH, Rizell M, Ericzon BG. Liver transplantation for cholangiocarcinoma: selection is essential for

acceptable results. *Scand J Gastroenterol.* 2011 Mar;46(3):370-5.

21. Jørgensen KK, Lindström L, Cvancarova M, Castedal M, Friman S, Schrumpf E, Foss A, Isoniemi H, Nordin A, Holte K, Rasmussen A, Bergquist A, Vatn MH, Boberg KM. Colorectal neoplasia in patients with primary sclerosing cholangitis undergoing liver transplantation: a Nordic multicenter study. *Scand J Gastroenterol.* 2012 Sep;47(8- 9):1021-9.

22. Jørgensen KK, Lindström L, Cvancarova M, Karlsen TH, Castedal M, Friman S, Schrumpf E, Foss A, Isoniemi H, Nordin A, Holte K, Rasmussen A, Bergquist A, Vatn MH, Boberg KM. Immunosuppression after liver transplantation for primary sclerosing cholangitis influences activity of inflammatory bowel disease. *Clin Gastroenterol Hepatol.* 2013 May;11(5):517-23

23. Fosby B, Melum E, Bjørø K, Bennet W, Rasmussen A, Andersen IM, Castedal M, Olausson M, Wibeck C, Gotlieb M, Gjertsen H, Toivonen L, Foss S, Makisalo H, Nordin A, Sanengen T, Bergquist A, Larsson ME, Soderdahl G, Nowak G, Boberg KM, Isoniemi H, Keiding S, Foss A, Line PD, Friman S, Schrumpf E, Ericzon BG, Höckerstedt K, Karlsen TH. Liver transplantation in the Nordic countries - An intention to treat and post-transplant analysis from The Nordic Liver Transplant Registry 1982-2013. *Scand J Gastroenterol.* 2015 Jun;50(6):797-808.

24. Thorsen T, Aandahl EM, Bennet W, Olausson M, Ericzon BG, Nowak G, Duraj F, Isoniemi H, Rasmussen A, Karlsen TH, Foss A. Transplantation With Livers From Deceased Donors Older Than 75 Years. *Transplantation.* 2015 Dec;99(12):2534-42

25. Åberg F, Gissler M, Karlsen TH, Ericzon BG, Foss A, Rasmussen A, Bennet W, Olausson M, Line PD, Nordin A, Bergquist A, Boberg KM, Castedal M, Pedersen CR, Isoniemi H. Differences in long-term survival among liver transplant recipients and the general population: a population-based Nordic study. *Hepatology.* 2015 Feb;61(2):668-77